

Lecture 10 Topics

- Bose-Einstein Statistics Applications
 - Blackbody radiation by photons
 - Laser: Light amplification by the stimulated emission of radiation

Three distributions

Distribution	Occupation index	Particles	Identical particles?	Spin	Distinguishable?	Exclusion principle?
Boltzman	$\frac{1}{Be^{E/k_B T}}$	Classical	Yes	Any spin	Yes	No
Bose-Einstein	$\frac{1}{Be^{E/k_B T} - 1}$	Bosons	Yes	0 or integer spin	No	No
Fermi-Dirac	$\frac{1}{Be^{E/k_B T} + 1}$	Fermions	Yes	1/2	No	Yes

Hadrons

TABLE 12.2 Commonly produced hadrons

Baryons	Mass (MeV/c ²)	Spin	Strange- ness	I, I_3	Lifetime, τ (or width \hbar/τ)	Mesons	Mass (MeV/c ²)	Spin	Strange- ness	I, I_3	Lifetime, τ (or width \hbar/τ)
p (uud)	938	$\frac{1}{2}$	0	$\frac{1}{2}, +\frac{1}{2}$	$>10^{32}$ yr	$\pi^+(u\bar{d})$	140	0	0	1, +1	2.6×10^{-8} s
n (udd)	940	$\frac{1}{2}$	0	$\frac{1}{2}, -\frac{1}{2}$	889 s	$\pi^0(u\bar{u} + d\bar{d})$	135	0	0	1, 0	8.4×10^{-17} s
Σ^+ (uus)	1189	$\frac{1}{2}$	-1	1, +1	8.0×10^{-11} s	$\pi^-(d\bar{u})$	140	0	0	1, -1	2.6×10^{-8} s
Σ^0 (uds)	1193	$\frac{1}{2}$	-1	1, 0	7.4×10^{-20} s	$K^+(u\bar{s})$	494	0	+1	$\frac{1}{2}, +\frac{1}{2}$	1.2×10^{-8} s
Λ^0 (uds)	1116	$\frac{1}{2}$	-1	0, 0	2.6×10^{-10} s	$K_S^0(d\bar{s}, s\bar{d})$	498	0	mix	$\frac{1}{2}, \text{mix}$	8.9×10^{-11} s
Σ^- (dds)	1197	$\frac{1}{2}$	-1	1, -1	1.5×10^{-10} s	$K_L^0(d\bar{s}, s\bar{d})$	498	0	mix	$\frac{1}{2}, \text{mix}$	5.2×10^{-8} s
Ξ^0 (uss)	1315	$\frac{1}{2}$	-2	$\frac{1}{2}, -\frac{1}{2}$	2.9×10^{-10} s	$K^-(s\bar{u})$	494	0	-1	$\frac{1}{2}, -\frac{1}{2}$	1.2×10^{-8} s
Ξ^- (dss)	1321	$\frac{1}{2}$	-2	$\frac{1}{2}, -\frac{1}{2}$	1.6×10^{-10} s	$\rho^+(u\bar{d})$	769	1	0	1, +1	151 MeV
Δ^{++} (uuu)	1232	$\frac{3}{2}$	0	$\frac{3}{2}, +\frac{3}{2}$	120 MeV	$\rho^0(u\bar{u} + d\bar{d})$	769	1	0	1, 0	151 MeV
Δ^+ (uud)	1232	$\frac{3}{2}$	0	$\frac{3}{2}, +\frac{1}{2}$	120 MeV	$\rho^-(d\bar{u})$	769	1	0	1, -1	151 MeV
Δ^0 (udd)	1232	$\frac{3}{2}$	0	$\frac{3}{2}, -\frac{1}{2}$	120 MeV	$K^{*+}(u\bar{s})$	892	1	+1	$\frac{1}{2}, +\frac{1}{2}$	50 MeV
Δ^- (ddd)	1232	$\frac{3}{2}$	0	$\frac{3}{2}, -\frac{3}{2}$	120 MeV	$K^{*0}(d\bar{s})$	896	1	+1	$\frac{1}{2}, -\frac{1}{2}$	51 MeV
Σ^{*+} (uus)	1383	$\frac{1}{2}$	-1	1, +1	~ 40 MeV	$\bar{K}^{*0}(s\bar{d})$	896	1	-1	$\frac{1}{2}, +\frac{1}{2}$	51 MeV
Σ^{*0} (uds)	1384	$\frac{1}{2}$	-1	1, 0	~ 40 MeV	$K^{*-}(s\bar{u})$	892	1	-1	$\frac{1}{2}, -\frac{1}{2}$	50 MeV
Σ^{*-} (dds)	1387	$\frac{1}{2}$	-1	1, -1	~ 40 MeV	Heavy mesons—containing quarks beyond the strange					
Ξ^{*0} (uss)	1532	$\frac{1}{2}$	-2	$\frac{1}{2}, +\frac{1}{2}$	~ 10 MeV	$J/\psi(c\bar{c})$	3100	1	0	0, 0	87 keV
Ξ^{*-} (dss)	1535	$\frac{1}{2}$	-2	$\frac{1}{2}, -\frac{1}{2}$	~ 10 MeV	$Y(bb)$	9460	1	0	0, 0	~ 50 keV
Ω^- (sss)	1672	$\frac{1}{2}$	-3	0, 0	8.2×10^{-11} s						

TABLE 12.1 Fundamental forces and particles

Force	Gravitation		Electroweak		Strong	Residual
Property	Mass/energy		Charge/weak charge		Color charge	
Strength	$\sim 10^{-39}$	$\sim 10^{-2}$	$\sim 10^{-6}$		1	
Range	$1/r^2$	$1/r^2$	10^{-3} fm		short	1 fm
Mediating Bosons	Graviton?	Photon, γ	W^+, W^-	Z^0	Gluon	π^\pm, π^0
Spin	2?	1	1	1	1	0
Mass	0?	$< 6 \times 10^{-22}$	80.4×10^3	91.2×10^3	< 10	140, 135
Charge	—	0	+1, -1	0	0	$\pm 1, 0$
Color charge	—	—	—	—	r, g, or b + $\bar{r}, \bar{g},$ or \bar{b}	Neutral


Quarks

Participants in gravitation, electroweak, and strong


	Spin	Mass	Charge	Color charge
Up, u	$\frac{1}{2}$	~ 5	$+\frac{2}{3}$	r, g, b
Down, d	$\frac{1}{2}$	~ 10	$-\frac{1}{3}$	r, g, b
Strange, s	$\frac{1}{2}$	~ 100	$-\frac{1}{3}$	r, g, b
Charm, c	$\frac{1}{2}$	$\sim 1.3 \times 10^3$	$+\frac{2}{3}$	r, g, b
Bottom, b	$\frac{1}{2}$	$\sim 4.5 \times 10^3$	$-\frac{1}{3}$	r, g, b
Top, t	$\frac{1}{2}$	$\sim 180 \times 10^3$	$+\frac{2}{3}$	r, g, b

Three distributions

Distribution	Occupation index	Particles	Identical particles?	Spin
Boltzman	$\frac{1}{Be^{E/k_B T}}$	Classical	Yes	Any spin
Bose-Einstein	$\frac{1}{Be^{E/k_B T} - 1}$	Bosons	Yes	0 or integer spin
Fermi-Dirac	$\frac{1}{Be^{E/k_B T} + 1}$	Fermions	Yes	1/2




Blackbody radiation



Stefan-Boltzmann law: The intensity of radiation $\propto T^4$

Blackbody radiation


Figure 6.24



$3.0 \times 10^{14} \text{ Hz}$

radiation $\propto T^4$


Blackbody radiation



Stefan-Boltzmann law: The intensity of radiation $\propto T^4$

Wien's law: The frequency associated with the maximum energy intensity is proportional to T


Blackbody radiation



Stefan-Boltzmann law: The intensity of radiation $\propto T^4$

Wien's law: The frequency associated with the maximum energy intensity is proportional to T

Blackbody radiation



Stefan-Boltzmann law: The intensity of radiation $\propto T^4$

Rayleigh-Jeans Approach

Consider a cubic cavity of $L \times L \times L$

The number of permissible states can be written as:

$$E_{(n_x, n_y, n_z)} = (n_x^2 + n_y^2 + n_z^2) \frac{\pi^2 \hbar^2}{2mL^2}$$

Rayleigh-Jeans Approach

Consider a cubic cavity of $L \times L \times L$

The number of permissible states can be written as:

$$n_x = \frac{2L}{\lambda} = 1, 2, 3, \dots$$

$$n_y = \frac{2L}{\lambda} = 1, 2, 3, \dots$$

$$n_z = \frac{2L}{\lambda} = 1, 2, 3, \dots$$

Rayleigh-Jeans Approach

Consider a cubic cavity of $L \times L \times L$

The number of permissible states can be written as:

$$n_x = \frac{2L}{\lambda} = 1, 2, 3, \dots$$

$$n_y = \frac{2L}{\lambda} = 1, 2, 3, \dots$$

$$n_z = \frac{2L}{\lambda} = 1, 2, 3, \dots$$

$$n^2 = n_x^2 + n_y^2 + n_z^2 = \left(\frac{2L}{\lambda}\right)^2 \text{ where } \begin{cases} n_x = 0, 1, 2 \dots \\ n_y = 0, 1, 2 \dots \\ n_z = 0, 1, 2 \dots \\ \text{except when all } n_x n_y n_z \text{ are zero} \end{cases}$$

Rayleigh-Jeans Approach

Consider a cubic cavity of $L \times L \times L$

The number of permissible states can be written as:

$$n_x = \frac{2L}{\lambda} = 1, 2, 3, \dots$$

$$n_y = \frac{2L}{\lambda} = 1, 2, 3, \dots$$

$$n_z = \frac{2L}{\lambda} = 1, 2, 3, \dots$$

$$n = \frac{2L}{\lambda} \rightarrow dn = -\frac{2L}{\lambda^2} d\lambda$$

$$n^2 = n_x^2 + n_y^2 + n_z^2 = \left(\frac{2L}{\lambda}\right)^2 \text{ where } \begin{cases} n_x = 0, 1, 2 \dots \\ n_y = 0, 1, 2 \dots \\ n_z = 0, 1, 2 \dots \\ \text{except when all } n_x, n_y, n_z \text{ are zero} \end{cases}$$

The number of permissible states can be written in terms of λ

$$N(n)dn = 2 \times \frac{1}{8} 4\pi n^2 dn = \pi n^2 dn =$$

Rayleigh-Jeans Approach

Consider a cubic cavity of $L \times L \times L$

The number of permissible states can be written as:

$$n_x = \frac{2L}{\lambda} = 1, 2, 3, \dots$$

$$n_y = \frac{2L}{\lambda} = 1, 2, 3, \dots$$

$$n_z = \frac{2L}{\lambda} = 1, 2, 3, \dots$$

$$n = \frac{2L}{\lambda} \rightarrow dn = -\frac{2L}{\lambda^2} d\lambda$$

$$n^2 = n_x^2 + n_y^2 + n_z^2 = \left(\frac{2L}{\lambda}\right)^2 \text{ where } \begin{cases} n_x = 0, 1, 2 \dots \\ n_y = 0, 1, 2 \dots \\ n_z = 0, 1, 2 \dots \\ \text{except when all } n_x n_y n_z \text{ are zero} \end{cases}$$

The number of permissible states can be written in terms of λ

$$N(n)dn = 2 \times \frac{1}{8} 4\pi n^2 dn = \pi n^2 dn = \pi \left(\frac{2L}{\lambda}\right)^2 \left(-\frac{2L}{\lambda^2} d\lambda\right) = -\frac{8\pi L^3 d\lambda}{\lambda^4} = -\frac{8\pi V d\lambda}{\lambda^4} \equiv -N(\lambda) d\lambda$$

Rayleigh-Jeans Approach

Since

$$\lambda = \frac{c}{\nu} \text{ and } d\lambda = -\frac{c}{\nu^2} d\nu$$

The number of permissible states in terms of ν

$$N(\nu)d\nu = -\frac{8\pi V d\lambda}{\lambda^4} = \boxed{\nu^3 d\nu}$$

Rayleigh-Jeans Approach

Since

$$\lambda = \frac{c}{\nu} \text{ and } d\lambda = -\frac{c}{\nu^2} d\nu$$

The number of permissible states in terms of ν

$$N(n)dn = -\frac{8\pi V d\lambda}{\lambda^4} = -\frac{8\pi V}{\left(\frac{c}{\nu}\right)^4} \left(-\frac{c}{\nu^2} d\nu\right) = \frac{8\pi V}{c^3} \nu^2 d\nu \equiv N(\nu) d\nu$$

the energy radiation rate can be expressed in terms of ν :

$$u(\nu) d\nu = k_B T N(\nu) d\nu =$$



Rayleigh-Jeans Approach

Since

$$\lambda = \frac{c}{\nu} \text{ and } d\lambda = -\frac{c}{\nu^2} d\nu$$

The number of permissible states in terms of ν

$$N(n)dn = -\frac{8\pi V d\lambda}{\lambda^4} = -\frac{8\pi V}{\left(\frac{c}{\nu}\right)^4} \left(-\frac{c}{\nu^2} d\nu\right) = \frac{8\pi V}{c^3} \nu^2 d\nu \equiv N(\nu) d\nu$$

the energy radiation rate can be expressed in terms of ν :

$$u(\nu) d\nu = k_B T N(\nu) d\nu = \frac{8\pi V k_B T}{c^3} \nu^2 d\nu$$


Rayleigh-Jeans Approach

Since

The number of permiss

$N(n)dn$

, the energy radiation



$$u(\nu) d\nu = k_B T N(\nu) d\nu = \frac{8\pi V k_B T}{c^3} \nu^2 d\nu$$

Planck's Hypothesis

$$E_n = nh\nu$$

$$\bar{E} = \sum E_n P(E_n) = \frac{\sum_n E_n e^{-\frac{E_n}{k_B T}}}{\sum_n e^{-\frac{E_n}{k_B T}}}$$

Planck's Hypothesis

$$E_n = nh\nu$$

$$\bar{E} = \sum E_n P(E_n) = \frac{\sum_n E_n e^{-\frac{E_n}{k_B T}}}{\sum_n e^{-\frac{E_n}{k_B T}}} = \frac{\sum_n nh\nu e^{-\frac{nh\nu}{k_B T}}}{\sum_n e^{-\frac{nh\nu}{k_B T}}}$$

Since $0 < e^{-\frac{nh\nu}{k_B T}} < 1$ and consider $e^{-\frac{nh\nu}{k_B T}} = x$

$$\bar{E} = \boxed{\phantom{\frac{nh\nu}{1 - e^{-\frac{nh\nu}{k_B T}}}}}$$

Planck's Hypothesis

$$E_n = nh\nu$$

$$\bar{E} = \sum E_n P(E_n) = \frac{\sum_n E_n e^{-\frac{E_n}{k_B T}}}{\sum_n e^{-\frac{E_n}{k_B T}}} = \frac{\sum_n nh\nu e^{-\frac{nh\nu}{k_B T}}}{\sum_n e^{-\frac{nh\nu}{k_B T}}}$$

Since $0 < e^{-\frac{h\nu}{k_B T}} < 1$ and consider $e^{-\frac{h\nu}{k_B T}} = x$

$$\bar{E} = h\nu \frac{\sum_n nx^n}{\sum_n x^n} :$$

Planck's Hypothesis

$$E_n = nh\nu$$

$$\bar{E} = \sum E_n P(E_n) = \frac{\sum_n E_n e^{-\frac{E_n}{k_B T}}}{\sum_n e^{-\frac{E_n}{k_B T}}} = \frac{\sum_n nh\nu e^{-\frac{nh\nu}{k_B T}}}{\sum_n e^{-\frac{nh\nu}{k_B T}}}$$

Since $0 < e^{-\frac{h\nu}{k_B T}} < 1$ and consider $e^{-\frac{h\nu}{k_B T}} = x$

$$\bar{E} = h\nu \frac{\sum_n nx^n}{\sum_n x^n} :$$

Using the following sums when $|x| < 1$

$$\sum_{n=0}^{\infty} x^n = \frac{1}{1-x} ; \sum_{n=0}^{\infty} nx^n = \frac{x}{(1-x)^2}$$

$$\bar{E} = h\nu \frac{\sum_n nx^n}{\sum_n x^n} =$$



Planck's Hypothesis

$$E_n = nh\nu$$

$$\bar{E} = \sum E_n P(E_n) = \frac{\sum_n E_n e^{-\frac{E_n}{k_B T}}}{\sum_n e^{-\frac{E_n}{k_B T}}} = \frac{\sum_n nh\nu e^{-\frac{nh\nu}{k_B T}}}{\sum_n e^{-\frac{nh\nu}{k_B T}}}$$

Since $0 < e^{-\frac{h\nu}{k_B T}} < 1$ and consider $e^{-\frac{h\nu}{k_B T}} = x$

$$\bar{E} = h\nu \frac{\sum_n nx^n}{\sum_n x^n} :$$

Using the following sums when $|x| < 1$

$$\sum_{n=0}^{\infty} x^n = \frac{1}{1-x} ; \sum_{n=0}^{\infty} nx^n = \frac{x}{(1-x)^2}$$

$$\bar{E} = h\nu \frac{\sum_n nx^n}{\sum_n x^n} = h\nu \frac{\frac{x}{(1-x)^2}}{\frac{1}{1-x}} = h\nu \frac{x}{1-x} = h\nu \frac{1}{x^{-1}-1} = h\nu \frac{1}{e^{\frac{h\nu}{k_B T}} - 1}$$

Planck's Hypothesis

$$E_n = nh\nu$$

$$\bar{E} = \sum E_n P(E_n) = \frac{\sum_n E_n e^{-\frac{E_n}{k_B T}}}{\sum_n e^{-\frac{E_n}{k_B T}}} = \frac{\sum_n nh\nu e^{-\frac{nh\nu}{k_B T}}}{\sum_n e^{-\frac{nh\nu}{k_B T}}}$$

Since $0 < e^{-\frac{nh\nu}{k_B T}} < 1$ and consider $e^{-\frac{nh\nu}{k_B T}} = x$

$$\bar{E} = h\nu \frac{\sum_n nx^n}{\sum_n x^n}$$

Using the following sums when $|x| < 1$

$$\sum_{n=0}^{\infty} x^n = \frac{1}{1-x}$$

$$\bar{E} = h\nu \frac{\sum_n nx^n}{\sum_n x^n} = h\nu \frac{\frac{x}{(1-x)^2}}{\frac{1}{1-x}} = h\nu \frac{x}{1-x} = h\nu \frac{1}{x^{-1}-1} = h\nu \frac{1}{e^{\frac{h\nu}{k_B T}} - 1}$$

Boltzman	$\frac{1}{Be^{E/k_B T}}$
Bose-Einstein	$\frac{1}{Be^{E/k_B T} - 1}$
Fermi-Dirac	$\frac{1}{Be^{E/k_B T} + 1}$

Energy

Planck's Hypothesis

The number of permissible states in terms of ν

$$E_n = nh\nu$$

$$N(n)dn = -\frac{8\pi V d\lambda}{\lambda^4} = -\frac{8\pi V}{\left(\frac{c}{\nu}\right)^4} \left(-\frac{c d\nu}{\nu^2}\right) = \frac{8\pi V}{c^3} \nu^2 d\nu$$

The number of permissible states can be rewritten in terms of energy

$$E = h\nu \text{ and } \nu = \frac{E}{h} \text{ and } d\nu = \frac{1}{h} dE$$

The total number of permissible states in terms of E would be

$$N(E)dE =$$



Planck's Hypothesis

The number of permissible states in terms of ν

$$E_n = nh\nu$$

$$N(n)dn = -\frac{8\pi V d\lambda}{\lambda^4} = -\frac{8\pi V}{\left(\frac{c}{\nu}\right)^4} \left(-\frac{c d\nu}{\nu^2}\right) = \frac{8\pi V}{c^3} \nu^2 d\nu$$

The number of permissible states can be rewritten in terms of energy

$$E = h\nu \text{ and } \nu = \frac{E}{h} \text{ and } d\nu = \frac{1}{h} dE$$

The total number of permissible states in terms of E would be

$$N(E)dE = \frac{8\pi V}{c^3 h^3} E^2 dE$$

Therefore, Density of States would be

$$D(E) = \square$$

Planck's Hypothesis

The number of permissible states in terms of ν

$$E_n = nh\nu$$

$$N(n)dn = -\frac{8\pi V d\lambda}{\lambda^4} = -\frac{8\pi V}{\left(\frac{c}{\nu}\right)^4} \left(-\frac{c d\nu}{\nu^2}\right) = \frac{8\pi V}{c^3} \nu^2 d\nu$$

The number of permissible states can be rewritten in terms of energy

$$E = h\nu \text{ and } \nu = \frac{E}{h} \text{ and } d\nu = \frac{1}{h} dE$$

The total number of permissible states in terms of E would be

$$N(E)dE = \frac{8\pi V}{c^3 h^3} E^2 dE$$

Therefore, Density of States would be

$$D(E) = \frac{8\pi V}{c^3 h^3} E^2$$

Planck's Hypothesis

$$\mathcal{N}(E) = \frac{1}{e^{\frac{h\nu}{k_B T} - 1}}$$

$$D(E) = \frac{8\pi V}{c^3 h^3} E^2$$

$$E = \int_0^{\infty} E \mathcal{N}(E) D(E) dE =$$

Planck's Hypothesis

$$\mathcal{N}(E) = \frac{1}{e^{\frac{h\nu}{k_B T} - 1}}$$

$$D(E) = \frac{8\pi V}{c^3 h^3} E^2$$

$$E = \int_0^\infty E \mathcal{N}(E) D(E) dE = \int_0^\infty \frac{E}{e^{E/k_B T} - 1} \left(\frac{8\pi V}{h^3 c^3} E^2 \right) dE$$

Use $E \equiv k_B T x$



Planck's Hypothesis

$$\mathcal{N}(E) = \frac{1}{e^{\frac{h\nu}{k_B T}} - 1}$$

$$D(E) = \frac{8\pi V}{c^3 h^3} E^2$$

$$E = \int_0^\infty E \mathcal{N}(E) D(E) dE = \int_0^\infty \frac{E}{e^{E/k_B T} - 1} \left(\frac{8\pi V}{h^3 c^3} E^2 \right) dE$$

Use $E \equiv k_B T x$

$$= \frac{8\pi V}{h^3 c^3} \int_0^\infty (k_B T)^4 \frac{x^3}{e^x - 1} dx$$



Planck's Hypothesis

$$\mathcal{N}(E) = \frac{1}{e^{\frac{h\nu}{k_B T}} - 1}$$

$$D(E) = \frac{8\pi V}{c^3 h^3} E^2$$

$$E = \int_0^\infty E \mathcal{N}(E) D(E) dE = \int_0^\infty \frac{E}{e^{E/k_B T} - 1} \left(\frac{8\pi V}{h^3 c^3} E^2 \right) dE$$

Use $E \equiv k_B T x$

$$= \frac{8\pi V}{h^3 c^3} \int_0^\infty (k_B T)^4 \frac{x^3}{e^x - 1} dx = \frac{8\pi V (k_B T)^4}{h^3 c^3} \frac{\pi^4}{15} = \frac{8V \pi^5 k_B^4}{h^3 c^3} T^4$$

$$\text{Since } \int_0^\infty \frac{x^3}{e^x - 1} dx = \frac{\pi^4}{15}$$

Planck's Hypothesis

$$E_n = nh\nu$$

$$D(E) = \frac{m^{3/2} V \sqrt{2}}{\pi^2 \hbar^3} \sqrt{E}$$

$$E = \int_0^\infty E N(E) D(E) dE = \int_0^\infty \frac{E}{e^{E/k_B T} - 1} \left(\frac{8\pi V}{h^3 c^3} E^2 \right) dE$$


Use $E \equiv k_B T x$

$$= \frac{8\pi V}{h^3 c^3} \int_0^\infty (k_B T)^4 \frac{x^3}{e^x - 1} dx = \frac{8\pi V (k_B T)^4}{h^3 c^3} \frac{\pi^4}{15} = \frac{8V \pi^5 k_B^4}{h^3 c^3} T^4$$

Stefan-Boltzmann's law

Planck's Hypothesis

$$dE = \frac{hv^3}{e^{hv/k_B T} - 1} \left(\frac{8\pi V}{c^3} \right) dv$$



Wien's Law

Planck's Hypothesis

$$dE = \frac{hv^3}{e^{hv/k_B T} - 1} \left(\frac{8\pi V}{c^3} \right) dv$$





Figure 6.24



LASER


Light Amplification by the Stimulated Emission of Radiation

Coherent Light

Figure 9.21 Coherent versus incoherent light.



Unidirectional
Monochromatic
Not in phase
Not coherent



Unidirectional
Monochromatic
In phase
Coherent

LASER

Light Amplification by the Stimulated Emission of Radiation


Coherent Light

- Moving in one direction
- Of a single wavelength
- In phase
- A lot of them

Figure 9.21 Coherent versus incoherent light.



Unidirectional
Monochromatic
Not in phase
Not coherent





Unidirectional
Monochromatic
In phase
Coherent

Einstein's theory

Spontaneous Emission:


$$R_{spo} = A_{spo} N_2$$



Einstein's theory


Spontaneous Emission:

$$R_{spo} = A_{spo} N_2$$



Absorption:


$$R_{abs} = B_{abs} N_1 Y(\Delta E)$$



Einstein's theory


Spontaneous Emission:

$$R_{spo} = A_{spo} N_2$$



Absorption:

$$R_{abs} = B_{abs} N_1 \rho(\Delta E)$$




The number of photons with the energy difference

Einstein's theory


Spontaneous Emission:

$$R_{spo} = A_{spo} N_2$$



Absorption:


$$R_{abs} = B_{abs} N_1 Y(\Delta E)$$



The number of photons with the energy difference

Stimulated Emission:


$$R_{sti} = B_{sti} N_2 Y(\Delta E)$$



Einstein's theory


Spontaneous Emission:

$$R_{spo} = A_{spo} N_2$$



Absorption:


$$R_{abs} = B_{abs} N_1 Y(\Delta E)$$



The number of photons with the energy difference

Stimulated Emission:

$$R_{sti} = B_{sti} N_2 Y(\Delta E)$$



Emission = Absorption

Einstein's theory

Emission = Absorption

$$R_{spo} + R_{sti} = R_{abs}$$

$$R_{spo} = A_{spo} N_2$$

$$R_{abs} = B_{abs} N_1 Y(\Delta E)$$

$$R_{sti} = B_{sti} N_2 Y(\Delta E)$$

Einstein's theory

Emission = Absorption

$$R_{spo} + R_{sti} = R_{abs}$$

$$A_{spo}N_2 + B_{sti}N_2Y(\Delta E) = B_{abs}N_1Y(\Delta E)$$

$$R_{spo} = A_{spo}N_2$$

$$R_{abs} = B_{abs}N_1Y(\Delta E)$$

$$R_{sti} = B_{sti}N_2Y(\Delta E)$$

Einstein's theory

Emission = Absorption

$$R_{spo} + R_{sti} = R_{abs}$$

$$A_{spo}N_2 + B_{sti}N_2Y(\Delta E) = B_{abs}N_1Y(\Delta E)$$

$$(B_{abs}N_1 - B_{sti}N_2)Y(\Delta E) = A_{spo}N_2$$

$Y(\Delta E)$:

$$R_{spo} = A_{spo}N_2$$

$$R_{abs} = B_{abs}N_1Y(\Delta E)$$

$$R_{sti} = B_{sti}N_2Y(\Delta E)$$

Einstein's theory

Emission = Absorption

$$R_{spo} + R_{sti} = R_{abs}$$

$$A_{spo}N_2 + B_{sti}N_2Y(\Delta E) = B_{abs}N_1Y(\Delta E)$$

$$(B_{abs}N_1 - B_{sti}N_2)Y(\Delta E) = A_{spo}N_2$$

$$Y(\Delta E) = \frac{A_{spo}N_2}{(B_{abs}N_1 - B_{sti}N_2)} = \frac{A_{spo}/B_{abs}}{\frac{N_1}{N_2} - \frac{B_{sti}}{B_{abs}}} =$$



$$\text{Since } N_1 \propto e^{-\frac{E_1}{k_B T}} \text{ and } N_2 \propto e^{-\frac{E_2}{k_B T}}$$

$$R_{spo} = A_{spo}N_2$$

$$R_{abs} = B_{abs}N_1Y(\Delta E)$$

$$R_{sti} = B_{sti}N_2Y(\Delta E)$$

Einstein's theory

Emission = Absorption

$$R_{spo} + R_{sti} = R_{abs}$$

$$A_{spo}N_2 + B_{sti}N_2Y(\Delta E) = B_{abs}N_1Y(\Delta E)$$

$$(B_{abs}N_1 - B_{sti}N_2Y)Y(\Delta E) = A_{spo}N_2$$

$$Y(\Delta E) = \frac{A_{spo}N_2}{(B_{abs}N_1 - B_{sti}N_2Y)} = \frac{A_{spo}/B_{abs}}{\frac{N_1}{N_2} - \frac{B_{sti}}{B_{abs}}} = \frac{A_{spo}/B_{abs}}{e^{\Delta E/k_B T} - \frac{B_{sti}}{B_{abs}}}$$

Einstein's theory

Emission = Absorption

$$R_{spo} + R_{sti} = R_{abs}$$

$$A_{spo}N_2 + B_{sti}N_2Y(\Delta E) = B_{abs}N_1Y(\Delta E)$$

$$(B_{abs}N_1 - B_{sti}N_2Y)Y(\Delta E) = A_{spo}N_2$$

$$Y(\Delta E) = \frac{A_{spo}N_2}{(B_{abs}N_1 - B_{sti}N_2Y)} = \frac{A_{spo}/B_{abs}}{\frac{N_1}{N_2} - \frac{B_{sti}}{B_{abs}}} = \frac{A_{spo}/B_{abs}}{e^{\Delta E/k_B T} - \frac{B_{sti}}{B_{abs}}}$$

= 1

If $N_1 = N_2$


Einstein's theory

Emission = Absorption

$$R_{spo} + R_{sti} = R_{abs}$$

$$A_{spo}N_2 + B_{sti}N_2Y(\Delta E) = B_{abs}N_1Y(\Delta E)$$

$$(B_{abs}N_1 - B_{sti}N_2Y)Y(\Delta E) = A_{spo}N_2$$




$$Y(\Delta E) = \frac{A_{spo}N_2}{(B_{abs}N_1 - B_{sti}N_2Y)} = \frac{A_{spo}/B_{abs}}{\frac{N_1}{N_2} - \frac{B_{sti}}{B_{abs}}} = \frac{A_{spo}/B_{abs}}{e^{\Delta E/k_B T} - \frac{B_{sti}}{B_{abs}}}$$

= 1


If $N_1 = N_2$

Three level Laser





Optical pumping



Three level Laser



Three level Laser



Three level Laser



Population Inversion!!!

Three level Laser

