

# Lecture 17

Thursday, March 08, 2012

## Quantum Oscillations (Landau levels; dHvA, SdH measurements)

We continue to consider the constant  $\vec{B}$  case without any  $\vec{E}$  field. As Landau found out, the  $\vec{B}$  field causes an additional quantization, which becomes very essential for quantum oscillation measurements (de Haas van Alphen effect etc.).

Recall the Bohr-Sommerfeld quantization condition from the semi-classical point of view:  $\oint d\vec{r} \cdot \vec{p} = (n + \gamma)h$  (Kittel has a typo for this equation; there is no  $2\pi$  if using  $h$ ). Here,  $\gamma$  is a sort of "greedy factor" that we use to fit the semi-classical theory even for the ground state. For this problem,  $\gamma = 1/2$  like in a simple Harmonic oscillator, but we won't need it.


We will consider the free electron case only, the generalization to the band case is straightforward.

Recall that the canonical momentum is given by  $m\vec{v} + \frac{q}{c}\vec{A} = \hbar\vec{k} + \frac{q}{c}\vec{A}$ . (In the case of a wave packet of Bloch states, this identity would be valid for each sub-wave packet (with each  $\vec{k}$  belonging to the same cell in the extended zone scheme) up to a constant  $\vec{G}$ ; that would be sufficient for the following derivation since the integral that we consider is independent of any constant offset in  $\vec{k}$ .)

We apply the Bohr-Sommerfeld quantization condition,  $\oint d\vec{r} \cdot \vec{p} = (n + \gamma)h$ , to the classical particle with the canonical momentum  $\vec{p} = m\vec{v} + \frac{q}{c}\vec{A} = \hbar\vec{k} + \frac{q}{c}\vec{A}$ .

$$\oint d\vec{r} \cdot \vec{p} = \hbar \oint d\vec{r} \cdot \vec{k} + \frac{q}{c} \oint d\vec{r} \cdot \vec{A} \quad \rightarrow \text{Stoke's theorem}$$

In the 2nd term:  $\oint d\vec{r} \cdot \vec{A} = \int d\vec{\sigma} \cdot \vec{\nabla} \times \vec{A} = \int d\vec{\sigma} \cdot \vec{B} = \Phi = \text{magnetic flux}$ .



$d\vec{r}$

For the 1st term, recall that  $d\vec{k} = \frac{q}{\hbar c} d\vec{r} \times \vec{B}$ , and so  $\vec{k} = \frac{q}{\hbar c} \vec{r} \times \vec{B} + \text{const}$  in the constant  $\vec{B}$  field. So, the 1st term:  $\hbar \oint d\vec{r} \cdot \vec{k} = \frac{q}{c} \oint d\vec{r} \cdot (\vec{r} \times \vec{B}) = \frac{q}{c} \vec{B} \cdot \oint d\vec{r} \times \vec{r} = -2 \frac{q}{c} B \Sigma$  where  $\Sigma$  is the area enclosed by the orbit. Since  $B\Sigma = \Phi$ , by collecting the first and the 2nd terms we get the following quantization condition:

$$-\frac{q}{c} \Phi = (n + \gamma)h$$

For the electron, we use  $q = -e$ , to get

$$\Phi_n = (n + \gamma) \frac{\hbar c}{e}$$

So, **the magnetic flux is quantized**. The flux quantum  $\Phi_0 = \frac{\hbar c}{e} = 4.14 \times 10^{-7}$  gauss cm<sup>2</sup>, and  $\Phi_n = (n + \gamma)\Phi_0$

In other words, an orbit area quantization occurs, both in  $\vec{r}$  space and in  $\vec{k}$  space, in this semi-classical picture. Let us call the area in  $\vec{r}$  space  $A_n$ , and the area in  $\vec{k}$  space  $S_n$ . The above flux quantization means that  $A_n B = (n + \gamma) \frac{\hbar c}{e}$ . And, since  $d\vec{k} = \frac{-e}{\hbar c} d\vec{r} \times \vec{B}$ , we have  $S_n = \left(\frac{eB}{\hbar c}\right)^2 A_n$ . Therefore, we have the following quantization for the orbital area in  $\vec{k}$  space.

$$S_n = 2\pi \frac{eB}{\hbar c} (n + \gamma)$$

This is the central formula that explains why the quantum oscillation phenomena occur.

Suppose one has a 2D electron gas. In the absence of the  $\vec{B}$  field, the allowed  $\vec{k}$  values form a practical continuum. As the  $\vec{B}$  field is turned on, though, not all  $\vec{k}$  values are accessible. Only those  $\vec{k}$  values on a circle, whose area corresponds to  $S_n$  are available.  $\vec{k}$  is not a good quantum number any more, but each circle represents a quantum number: this is the **Landau level**, which is just the index of concentric circles in  $\vec{k}$  space, or equivalently in  $\vec{r}$  space. [In some conventions, the Landau level starts from  $n = 1$ , in which case  $\gamma = -1/2$ , instead of  $1/2$ . Here we will use  $n = 0, 1, 2, \dots$  and  $\gamma = 1/2$ .]

It is convenient to define a "magnetic length scale"  $l_B = \sqrt{\frac{\hbar c}{eB}}$  to express the above results as:


$$S_n = \frac{2\pi}{l_B^2}(n + \gamma), A_n = l_B^4 S_n, A_n = 2\pi l_B^2(n + \gamma)$$

With  $\gamma = 1/2$ , and  $n = 0, 1, 2, \dots$ , one sees that  $l_B$  ( $1/l_B$ ) is the radius of the first Landau orbit in  $\vec{r}$  ( $\vec{k}$ ) space.  $l_B = \frac{257}{\sqrt{B}} \text{ \AA}$ , if  $B$  was in unit of Tesla. So, in accessible  $B$  field range,  $l_B$  would be a few tens of  $\text{\AA}$  or more, or  $2\pi l_B \gtrsim 100 \text{ \AA}$ . In terms of flux quantum  $\Phi_0$ ,  $l_B = \sqrt{\frac{\Phi_0}{2\pi B}}$ : i.e.,  $\pi l_B^2$  corresponds to half the flux quantum.

Landau levels have a degeneracy that increases with  $B$ . From the above equation,  $\Delta S_n = \frac{2\pi}{l_B^2}$ . It is reasonable to conjecture that the degeneracy of each Landau level is just this divided by the usual  $(2\pi)^2/A_S$ , where  $A_S$  is the area of the two dimensional crystal (or the crosssectional area of the three dimensional crystal). Not surprisingly, Quantum Mechanics proves this conjecture to be true. So, the Landau degeneracy is  $D = \frac{1}{2\pi} \frac{A_S}{l_B^2} = \frac{BA_S}{\Phi_0} = \Phi_{\text{tot}}/\Phi_0$ , where  $\Phi_{\text{tot}}$  is the magnetic flux through the sample. Including the spin degeneracy, we get

$$D = 2\Phi_{\text{tot}}/\Phi_0$$

The physics of Landau levels underlies all quantum oscillation measurements.



In the current semi-classical picture, the above picture describes what happens as we turn on the  $\vec{B}$  field. Without the  $\vec{B}$  field, the states up to  $\vec{k} = \vec{k}_F$  are occupied. With the  $\vec{B}$  field, however, only those circles of  $\vec{k}$  values are allowed (keeping in mind, of course, that, in the quantum mechanical sense, there would be only dense and sparse regions of  $\vec{k}$  space, in terms of the probability density). For a

given  $\vec{B}$  field, there would be an outermost orbit which is partially-occupied or un-occupied while all smaller orbits are fully occupied. As  $\vec{B}$  field is increased, the orbits will become large, while at the same time their Landau degeneracy increases. Therefore, the outermost orbital will be emptied gradually, until it becomes completely empty and then the number of occupied orbitals decrease by 1. This is the origin of all quantum oscillation measurements in the  $\vec{B}$  field.

How does this make it possible to measure the Fermi surface? Note that from the above diagram the emptying of the outermost orbital will come when the area of the outermost orbital is equal to the area of the Fermi surface plus  $\frac{2\pi}{l_B^2} \gamma$ .

Thus,  $S_n = \frac{2\pi}{l_B^2} (n + \gamma) = A_{FS} + \frac{2\pi}{l_B^2} \gamma$  defines the condition for the "jumps" where  $A_{FS}$  is the area of the Fermi surface.  $\frac{4\pi^2 B}{\Phi_0} n = A_{FS}$ , or  $\frac{1}{B} = \frac{4\pi^2}{\Phi_0 A_{FS}} n$ . This is the origin of oscillations in physical quantities as a function of  $\frac{1}{B}$ . The period in  $\frac{1}{B}$  is given by  $\frac{4\pi^2}{\Phi_0 A_{FS}}$ .

$$\Delta\left(\frac{1}{B}\right) = \frac{4\pi^2}{\Phi_0} \frac{1}{A_{FS}} = \frac{2\pi e}{\hbar c} \frac{1}{A_{FS}}$$


So, by measuring the period,  $A_{FS}$  can be measured!

It is a well-known QM result that each Landau level has the energy  $(n + \frac{1}{2}) \hbar \omega_c$ , where  $\omega_c = \frac{eB}{mc} = \frac{\hbar}{m} \frac{1}{l_B^2}$  is the cyclotron frequency. In the above semi-classical model, this can be obtained as  $\frac{p^2}{2m} = \frac{\hbar^2}{2m} \frac{S_n}{\pi} = \frac{\hbar^2}{m} \frac{1}{l_B^2} (n + \gamma) = \hbar \omega_c (n + \gamma)$ .

In order for the quantum oscillation measurements to succeed, it is necessary that the sample is pure (the mean free path  $l \gg l_B$ ) and the temperature is very low ( $k_B T \ll \hbar \omega_c$ ).

So far, we have considered only a two dimensional case. What about three dimensions? Here the so called "one dimensional density of states (1D DOS) effect" comes in. For a given dispersion  $\epsilon(\vec{k})$ , consider the 1D density of states along the  $z$  direction, which is defined as the direction of the  $\vec{B}$  field. For a given energy and fixed  $k_x, k_y$  values, one can think about the 1D DOS, in the sense of the available number of  $k_z$  values. Thus, 1D DOS  $\propto \frac{1}{|\frac{\partial \epsilon(\vec{k})}{\partial k_z}|} \propto \frac{1}{|v_{g,z}|}$ . The

importance of this 1D DOS is that when it diverges, that  $k_z$  value will singularly contribute the response of the system.



This is illustrated in the above diagram. While quantum oscillation phenomena happen at any fixed  $k_z$  value, the three **extremal Fermi surfaces**  $\left(\frac{\partial \epsilon(\vec{k})}{\partial k_z} = 0\right)$  shown in the diagram are the only ones that show up in the actual measurements, since they dominate the response function, because the 1D DOS is infinite at those  $k_z$  values. In the case illustrated here, the two or three oscillation frequencies will beat.

The oscillation in  $1/B$  can be measured in a variety of physical quantities (conductivity, magnetization, specific heat etc.). For instance, when the magnetization (or the magnetic susceptibility) is measured and plotted as a function of  $1/B$ , an oscillatory behavior can be observed. The period of that oscillation  $\left(\frac{2\pi e}{\hbar c} \frac{1}{A_{FS}}\right)$  gives information about the area of the extremal FS. This effect is called the **de Hass van Alphen (dHvA)** effect. When the conductivity shows an oscillatory behavior, then it is called the **Schubnikov de Hass (SdH)** effect. These are very important techniques in probing the Fermi surface.