

# Notes for Lecture 11

## Phonons and photons

We have established the quantum mechanical formalism. With this formalism, we will explore some important quantum particles in the next few lectures. We assume that they are free, while keeping in mind that they must interact to some extent since otherwise the equilibrium cannot be established.

### 11.1 Phonons

Phonons refer to quantized lattice vibrations. They are elementary but their applications in practice and in conceptual matters are quite broad. So, it is important to know them well.

#### 11.1.1 Normal modes

Before we can discuss phonons, we must discuss the *normal modes* of the coupled oscillator problem in classical mechanics. If you are not familiar with this problem, you might want to read up on your old note in classical mechanics<sup>1</sup>.

The grand result is the following: for a three dimensional solid that consists of  $N$  atoms, there must be  $3N$  normal modes, which can be thought of as completely independent simple harmonic oscillators within the so-called *harmonic approximation*. The harmonic approximation means that the inter-atomic potential is approximated to the second order in displacements with respect to the equilibrium, just as in the

---

<sup>1</sup> My teaching web site for 105 may be of help.

usual “Hooke’s law” approximation. In general, it can be shown that the Lagrangian can be expressed as a quadratic form involving two real symmetric matrices, one for the kinetic energy and the other for the potential energy; it can then be proved formally that the problem can always be diagonalized to yield exactly  $3N$  completely independent simple harmonic oscillators. These independent modes are what we refer to as “normal modes.”

So, it means that the Hamiltonian of *any* stable object (be it a molecule or a solid) can be written as

$$H = \sum_{i=1}^{3N} \left( \frac{p_{\eta,i}^2}{2m_i} + \frac{1}{2} m_i \omega_i^2 \eta_i^2 \right) \quad (11.1)$$

where  $\eta_i$  is the normal mode coordinate,  $p_{\eta,i}$  is its conjugate momentum, and  $m_i$  is the effective mass, and  $\omega_i$  the frequency, of each normal mode, as long as the displacements of the atoms are not too large<sup>2</sup>.

Quantizing the above equation we get

$$H = \sum_{i=1}^{3N} \left( a_i^\dagger a_i + \frac{1}{2} \right) \hbar \omega_i \quad (11.2)$$

where  $a_i, a_i^\dagger$  are the lowering and raising operators that can be formed by “taking the square root of the Hamiltonian,” so to speak

$$a_i = \sqrt{\frac{1}{2\hbar\omega_i}} \left( \sqrt{m_i}\omega_i\eta_i + i \frac{p_{\eta,i}}{\sqrt{m_i}} \right) \quad (11.3)$$

$$a_i^\dagger = \sqrt{\frac{1}{2\hbar\omega_i}} \left( \sqrt{m_i}\omega_i\eta_i - i \frac{p_{\eta,i}}{\sqrt{m_i}} \right) \quad (11.4)$$

These two operators satisfy the following fundamental relation

$$[a_i, a_j^\dagger] = \delta_{i,j} \quad (11.5)$$

While elementary quantum mechanics books refer to these operators as simply lowering and raising operators, they are, in fact, **the annihilation, or the destruction, operator and the creation operator for a boson**. What is this boson in this case? It is a particle in term of which we can describe how vigorously a string vibrates. Quantum mechanically, we describe the high energy state of a vibrating string as the state in which there are lots of lots of particles – lots of vibration quanta. In

---

<sup>2</sup> If the displacements are significant, then the normal modes are independent only in an approximate sense. Many interesting non-linear effects occur, which involve their modification and mixing.

other words, lots of *phonons*. By the same token, this same mathematical formalism can describe photons as well. Photons are much like phonons, as we shall see, except that they are related to light wave not sound wave. When you are looking at a bright object, you are being hit by lots and lots of photons. So, a bright light corresponds to lots of photons. The precise meaning of “lots of photons,” or “lots of phonons” is simply that  $n_i = a_i^\dagger a_i$  is a high number<sup>3</sup>.

What are these indices  $i, j$ ? They are simply labels for vibrational modes. However, note that each vibrational mode is an eigenstate of a Lagrangian, which could well have certain symmetries. For instance, in a crystal, the Lagrangian will have crystal translation symmetry as well as certain crystal rotational symmetry and others. It then follows that the normal modes can be taken as eigenstates of the corresponding translation, rotation or other operation. Thus, in general, the index  $i$  can be taken as a symmetry label. Notably, in a crystal, it can be taken as, among other things, the “crystal momentum,”  $\vec{k}$ . The full explanation of the crystal momentum is out of the scope of this course<sup>4</sup>. Here, a one dimensional crystal example suffices. Consider a one crystal consisting with the lattice constant  $a$ . This means that the crystal consists of a unit that is repeated by the translation by  $a$  indefinitely. It turns out that this view of a crystal as “something that repeats” is not just applicable in the real space, but applicable just as well in the wave vector space. In the wave vector space, any wave number  $k$  is equivalent to  $k + 2\pi/a$  in this one dimensional crystal. This defines the notion of the crystal momentum. It ( $\hbar\vec{k}$ ) is just like the momentum, except that its value itself is periodic. In this one dimensional example,  $\hbar k$  and  $\hbar(k + 2\pi/a)$  are completely equivalent, due to these wave vectors applicable to waves in crystal. In addition to the crystal momentum, the label  $i$  can have other symmetry labels, and so  $i$  is a collective index. In particular, note that it will in general, in dimensions higher than 1, include the polarization label. A similar consideration applies to photons, as well:  $i$  is a collective index for the momentum  $\vec{k}$  and the polarization.

**Phonon is a vibration quantum, and  $n_i = a_i^\dagger a_i$  corresponds to the number of phonons.** The quantum number  $i$  can be wave vector, polarization, or some other branch index (e.g., Debye phonon branch, Einstein phonon branch). As we shall see below, there is a one-to-one mathematical analogy between photon and low energy phonon.

---

<sup>3</sup> The commutation relation of Eq. 11.5 is the canonical commutation relation for bosons. Fermion creation and destruction operators satisfy  $\{a_i, a_j^\dagger\} = \delta_{i,j}$ , where  $\{a, b\} \equiv ab + ba$  is the *anti-commutator*. The formulation of the many body problem with these creation/destruction operators is referred to as the *second quantization*, which is beyond the scope of this course. However, any theoretically inclined person must study it at some point in a physics graduate school. Essentially, the second quantization is the scheme in which the wave function symmetry is kept track of most efficiently.

<sup>4</sup> My lecture notes for physics 155 may be of help.

### 11.1.2 Dispersion relation

With  $i$  in  $\omega_i$  being identified as a symmetry label, which include the crystal momentum and the polarization, one can ask what is the relationship between say  $\vec{k}$  (crystal momentum) and  $\omega$  for a given polarization?

Such a relationship is called the *dispersion relation*

$$\omega = \omega(\vec{k}) \quad (11.6)$$

As  $\hbar\vec{k}$  is the momentum and  $\hbar\omega$  is the energy, the dispersion relation can be viewed as defining the relation between the energy and the momentum.

### 11.1.3 Einstein phonons

This seemingly crude model has survived to this day, while it had not achieved its original goal in full, namely explaining the low temperature heat capacity of the diamond (and other solids).

The Einstein model simply assumes that all normal modes inside a solid is a single frequency mode.

$$\omega(\vec{k}) = \omega_E \quad (11.7)$$

You might ask how is this possible? Indeed, it may seem quite unlikely. However, for a crystal in which the repeating unit consists of more than one atom, there are phonon modes called “optical phonons.” The Einstein model has survived as the simplest model to describe these optical phonons.

Now, given the fact that the Einstein model considers  $3N$  independent oscillators of the same frequency, it follows that the thermodynamics of Einstein phonons is exactly the same as what has been described in Section 8.5, except that we have to apply the following transformations:  $\omega \rightarrow \omega_E$ ,  $Z \rightarrow Z^{3N}$ ,  $E \rightarrow 3NE$ ,  $C_V \rightarrow 3NC_V$  etc.

Therefore, the following qualitative behavior, much like the one oscillator problem, persists. (1) At high temperature ( $k_B T \gg \hbar\omega_E$ ), the system behaves classically, and obeys the equipartition theorem and, thus, the Du-Long Petit law  $C_V = 3Nk_B$ . (2) At low temperature ( $k_B T \ll \hbar\omega_E$ ), the system behaves quantum mechanically, and shows an activated behavior: e.g. the temperature dependence of  $C_V$  is dominated by  $\exp\left(-\frac{\hbar\omega_E}{k_B T}\right)$ .

While this model was originally suggested for explaining the heat capacity of the diamond, the behavior (2) is unlike what is actually observed. The actually observed

behavior is  $C_V \propto T^3$ , the Debye- $T^3$  law, which needs to be explained with the help of universal<sup>5</sup> acoustic phonons (sound waves).

### 11.1.4 Debye phonons

In the Debye model, the photon dispersion relation is assumed to be

$$\omega(\vec{k}) = vk \qquad k \equiv |\vec{k}| \qquad (11.8)$$

Note that this becomes exactly the dispersion relation for photons in vacuum, if one substitutes  $c$  for  $v$ .

The phonons that obey the above dispersion relation are called Debye phonons. In solids, sound waves propagate as waves that satisfy the above dispersion relation at long wave length. So, sound waves, or acoustic phonons, are Debye phonons at long wave lengths. For them,  $v$  is about  $10^{-5}$  of  $c$ .

While the Einstein model failed to explain the heat capacity of the diamond, the Debye model succeeded in explaining it, and as the result Debye received a big prize. Why the difference?

The difference lies in the fact that **acoustic phonons are universal**. The reason is deep. Consider a crystal place in vacuum. Now consider a wave (in other words, a particle) existing inside the crystal. Then, it is clear that from the point of view of that wave, the continuous translation symmetry of the vacuum is broken. By the Goldstone theorem, this situation allows the so-called Goldstone mode to exist. The Goldstone mode is a bosonic wave, whose energy goes to zero as the momentum goes to zero. More intuitively, the Goldstone mode can be thought of as the oscillation of the system that locally exploits the continuous symmetry that used to exist. The acoustic phonon case exemplifies such a mode perfectly. “Exploiting the continuous symmetry that used to exist” means, in this case, all atoms moving in one direction. The translational symmetry of the vacuum means that there is no restoring force for such a motion. Now, consider a sound wave, and make the wave length grow larger and larger. You have a wave whose local atomic displacements are just those of a translation, which does not cause any restoring force. Thus, a sound wave frequency goes to zero as  $k \rightarrow 0$ . This is the precise sense in which the sound wave is a Goldstone mode associated with the broken translational (and rotational) symmetry state of a crystal.

Note that this discussion assumes that the atomic interactions are local. If there is a long-range interaction, then this argument does not hold. Also, it happens that

---

<sup>5</sup> In contrast, optical phonons are not universal, as they do not exist in a monatomic crystal.

for typical short-ranged atomic interactions, the sound wave frequency goes to zero exactly as the above dispersion, i.e.  $\omega \propto k$ .

Here are some key steps for the derivation of the Debye  $T^3$  law. Throughout this derivation, it is critical to keep in mind that, for *each* oscillator mode defined by  $\omega$ , what we did for the Einstein mode applies.

Using the periodic boundary condition for wave in a box, we get that each  $\vec{k}$  value “owns” a volume element of

$$\Delta k = \frac{8\pi^3}{V} \quad (11.9)$$

By considering a sphere with radius  $k$ , dividing the volume of the radius by  $\Delta k$ , we get the number of phonon modes,  $N_m$ , in that volume.  $N_m = \frac{4\pi k^3}{3} / \Delta k$ . Expressing  $k$  in terms of  $\omega$  using the dispersion relation, and then taking the derivative, we get the density of states

$$D(\omega) = \frac{3V\omega^2}{2\pi^2 v^3} \quad (11.10)$$

where the 3 in the numerator takes into account the three polarizations per  $\vec{k}$ . We have the particle sum rule

$$\int_0^{\omega_D} d\omega D(\omega) = 3N \quad (11.11)$$

with  $N$  given by the number of atoms. From this, one gets

$$\omega_D = v(6\pi^2 n)^{1/3} \quad (11.12)$$

and

$$D(\omega) = \frac{9N\omega^2}{\omega_D^3} \quad (11.13)$$

The frequency scale  $\omega_D$  (Debye frequency) leads to the definition of other scales

$$\theta_D = \frac{\hbar\omega_D}{k_B} \quad \text{Debye temperature} \quad (11.14)$$

$$k_D = \frac{\hbar\omega_D}{v} \quad \text{Debye wave vector} \quad (11.15)$$

By carrying out the integral  $E = E_0 + \int_0^{\omega_D} d\omega D(\omega) n_P(\hbar\omega) \hbar\omega$  where  $n_P$  is the Planck distribution function (Eq. 8.60) we get

$$E = E_0 + 9Nk_B T \left( \frac{T}{\theta_D} \right)^3 \int_0^{\frac{\theta_D}{T}} dx \frac{x^3}{e^x - 1} \quad (11.16)$$

This leads to

$$C_V \approx \begin{cases} 3Nk_B & \text{if } T \gg \theta_D \\ \frac{12\pi^4}{5} Nk_B \left(\frac{T}{\theta_D}\right)^3 & \text{if } T \ll \theta_D, \text{ Debye } T^3 \text{ law} \end{cases} \quad (11.17)$$

It is important to be able to understand these results, qualitatively. When  $T \ll \theta_D$ , only phonons with  $\omega \lesssim \omega_{max} = k_B T / \hbar$  are excited substantially. These are “classical phonons,” which contribute dominantly to the thermodynamics. The maximum wave vector for these classical phonons is  $k_{max} = \omega_{max} / v \propto T$ . Thus, the number of all classical phonon modes is  $\propto k_{max}^3 \propto T^3$ . The total energy is obtained by multiplying this by  $k_B T$ , the equipartition energy per mode. So,  $E - E_0 \propto T^4$ . By dimensional argument and the extensivity, we must have  $E - E_0 = a N k_B T \left(\frac{T}{\theta_D}\right)^3$ , where  $a$  is a dimensionless constant. Then,  $C_V = 4a N k_B \left(\frac{T}{\theta_D}\right)^3$ . Thus, we get the correct solution from this qualitative argument up to a numerical factor.

## 11.2 Photons

Photons are just like Debye phonons, except for a few differences. (1) The speed in vacuum is given by the constant  $c$ . (2) There are only two polarizations in vacuum. (3) We do not know what medium, if any, corresponds to photon’s vibration. (4) There is no fixed number of photon *modes*. In contrast, there are only  $ND$  phonon modes, for  $N$  atoms in  $D$  dimensions.

By re-using the results of the Debye phonon problem, or simply re-doing it with the density of states and so on, one gets

$$E = E_0 + V \frac{\pi^2}{15} \left(\frac{k_B T}{\hbar c}\right)^3 k_B T \quad (11.18)$$

where  $E_0$  is the zero point energy from all photons<sup>6</sup>. Now, from the energy, the pressure can be obtained very simply

$$P = \frac{1}{3} \frac{E}{V} \quad (11.19)$$

The reason is because in the current problem, the only volume dependence in the partition function is through the dispersion relation  $\omega(\vec{k}) = ck$  where  $k \propto V^{-1/3}$ . And,

---

<sup>6</sup> So,  $E_0 = \infty$ , even for a finite volume ( $E_0/V = \infty$ ). However, we keep it here, since its change due to geometry is still a meaningful quantity. The van der Waals force or the Casimir force is the result of such a change.

so for a microstate energy  $E_{\mu}$ , we have  $E_{\mu} \propto V^{-1/3}$ , and thus,

$$\frac{\partial E_{\mu}}{\partial V} = -\frac{1}{3} \frac{E_{\mu}}{V}$$

So, it follows that

$$\begin{aligned} \left(\frac{\partial Z}{\partial V}\right)_T &= \sum_{\mu} \frac{\partial \exp(-\beta E_{\mu})}{\partial V} \\ &= \frac{\beta}{3} \sum_{\mu} \frac{E_{\mu}}{V} \exp(-\beta E_{\mu}) \\ &= \frac{\beta}{3} \frac{E}{V} \end{aligned}$$

Using this and the fact that  $P = -\frac{\partial F}{\partial V} = k_B T \frac{\partial \log Z}{\partial V}$ , the above result that  $P = \frac{1}{3} \frac{E}{V}$  follows easily. Note that this result applies to the whole  $E$ , not just the thermal part  $E - E_0$ .

Now, let us ask the question – suppose we have a cavity filled with light, and we prepare a hole<sup>7</sup> on one of the walls. What is **the rate of effusion** of photons, i.e., the number of photons coming out per unit time per unit area? Let us define this rate<sup>8</sup> as  $R_N$ .

$$R_N = \frac{1}{V} \int dN_m n_P(\omega) c \cos \theta \quad (11.20)$$

where  $dN$  is the number of photon modes, and we have taken the surface normal to be the  $z$  axis, and  $\theta$  is the polar angle in the usual spherical coordinate convention. In the above expression the integral is over only those modes for which  $0 < \theta < \pi/2$ , since otherwise the photon will not escape. Note that  $dN_m = D(\omega)d\omega$ , if we were doing an isotropic integral. Here, we are not. In fact,  $dN_m$  must involve  $d\omega$  and  $d(\cos \theta)$  in the current case. Still, we can make a clever use of this, since,  $dN_m = D(\omega)d\omega$  must result *if* we were allowed to do the  $d(\cos \theta)$  integral without worrying about the integrand. So, we must have

$$dN_m = D(\omega)d\omega \frac{d(\cos \theta)}{2}. \quad (11.21)$$

Plugging this into the above expression for  $R$ , and effecting the integral, while remembering the integration range for  $\cos \theta$  is only from 0 to 1, not from -1 to 1, we

---

<sup>7</sup> For the flow of particles to be an effusion, rather than a hydrodynamic flow, as we assume here, the hole must be small, compared to the mean free path. This condition is satisfied by photons, automatically, since they do not directly interact with one another.

<sup>8</sup> Note that this rate is physically a “flux.”

get

$$R_N = \frac{1}{4V} \int d\omega D(\omega) n_P(\omega) c = \frac{nc}{4} \quad (11.22)$$

$$n \equiv \frac{N}{V} \quad \text{number density of photons}$$

Note that in the above derivation, we used the density of states and the distribution function without needing any details about them. Thus, the above result for  $R_N$  can be generalized to *any* particle as<sup>9</sup>

$$R_N = \frac{n}{4} \overline{\langle v \rangle} \quad n: \text{ number density} \quad (11.23)$$

where  $v$  is the speed<sup>10</sup>. Similarly, one can define the rate of effusion for energy,  $R_E$ , for which we merely need to multiply the above integrand by  $\hbar\omega$ .

$$R_E = \frac{u}{4} \overline{\langle v \rangle} \quad \text{rate of effusion for energy} \quad (11.24)$$

$$u \equiv \frac{E}{V} \quad \text{energy density} \quad (11.25)$$

Again, this expression is valid for any particle. For photons, the speed is constant,  $c$ , and using<sup>11</sup>  $E - E_0$  in Eq. 11.18 for  $E$ , we get

$$R_E = \frac{c}{4} \frac{E - E_0}{V} = \sigma T^4 \quad \sigma = \frac{\pi^2 k_B^4}{60 \hbar^3 c^2} \quad (11.26)$$

which is the celebrated **Stefan's law**, with the Stefan's constant,  $\sigma = 5.670 \times 10^{-8} \text{ Wm}^{-2}\text{K}^{-4}$ .

<sup>9</sup> You are reminded that the quantum mechanical ensemble average is given the symbol  $\overline{\langle O \rangle}$  for an operator  $O$ . See Eq. 10.19.

<sup>10</sup> The speed is given by  $c$  (massless) or  $\hbar k/m$  (massive), for each free particle considered here.

<sup>11</sup> Since  $E_0$  is everywhere, it is not transported by effusing particles. More precisely speaking, particles are defined only with respect to the vacuum, and so particles carry the energy excitation energy, not the vacuum energy (i.e., the ground state energy).